



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

*Mortuary statistics from the State of Sao Paulo.*

*Sao Paulo* (capital of State).—Week ended January 8, 1905. Total deaths, 115. Stillbirths, 9. Causes of death, variola, 1; measles, 1; whooping cough, 2; enteric fever, 2; dysentery, 1; malarial fevers, 4; pulmonary tuberculosis, 3; syphilis, 2; other general diseases, 1; diseases of the nervous system, 5; of the circulatory system, 20; of the respiratory system, 10; of the digestive system, 41; of the urinary system, 1; puerperal septicæmia, 1; congenital debility, 4; senile debility, 1; violence, except suicide, 2; diseases ill defined, 13.

By ages, less than 2 years, 67; 2 to 5 years, 4; 5 to 10 years, 3; 10 to 20 years, 4; 20 to 50 years, 19; more than 50 years of age, 17, and age unknown, 1. Natives, 95; foreigners, 19; nationality unknown, 1.

*Santos*.—Week ended January 8, 1905. Total deaths, 29. Causes of death, variola, 1; whooping cough, 1; malarial fevers, 2; tuberculosis, 3; diseases of the nervous system, 2; of the circulatory system, 4; of the respiratory system, 4; of the digestive system, 6; congenital debility, 1; senile debility, 1; violence, except suicide, 2; and diseases ill defined, 2. Natives, 23; foreigners, 6.

*Campinas*.—Week ended January 8, 1905. Total deaths, 28. Causes of death: Whooping cough, 2; diphtheria, 1; grippe, 1; dysentery, 1; tuberculosis, 3; septicæmia, 1; diseases of nervous system, 2; of the circulatory system, 1; of the respiratory system, 3; of the digestive system, 5; congenital debility, 2; violence, except suicide, 1; and diseases ill defined, 5. Natives, 20; foreigners, 8. Daily average number of deaths, 4.

*Mortuary report of Rio de Janeiro, week ended January 29, 1905.*

During the week there were, in all, 299 deaths. Of this number, 1 was caused by yellow fever, with 10 new cases. Last week there were 4 cases and this week 10 cases. This seems to show that there may be a recrudescence of fever here this season. Still, it may be only a temporary disturbance.

I have not observed many mosquitoes lately, though the weather has been very hot, followed by heavy rains, which, of course, is a favorable condition for the breeding of mosquitoes.

Plague had 11 new cases and 1 death, leaving at the end of the week 51 cases under treatment. Variola caused 14 deaths, with 33 new cases, and at the close of the week there were 76 cases in Sao Sebastiao Hospital. Of the cases of yellow fever, 5 are in hospital and the remainder are being treated at their homes.

Other causes of death were: Measles, 5; grippe, 12; enteric fever, 1; dysentery, 4; beriberi, 4; leprosy, 1; erysipelas, 2; malarial fevers, 10 (last week 7, week before that 3—also showing increase, probably due to mosquitoes); tuberculosis, pulmonary, 48; other forms of tuberculosis, 3; septicaemia, 2; syphilis, 2; cancer, 5; other general diseases, 1; diseases of the nervous system, 31; of the circulatory system, 45; of the respiratory system, 23; of the digestive system, 45; of the urinary system, 11; congenital debility, 9; senile debility, 9; violence, except suicide, 6; suicide, 3; diseases ill defined, 1.

By ages, less than 1 year old, 66; 1 to 5 years, 21; 5 to 10, 12; 10 to 20, 17; 20 to 30, 35; 30 to 40, 32; 40 to 50, 35; 50 to 60, 27; more than 60 years old, 54.